Biotechnic & Histochemistry ›
Volume 91, 2016 - Issue 7

376 Views | 7 CrossRef citations to date | 1 Altmetric

Original Articles

Effects of long-term exposure to 900 megahertz electromagnetic field on heart morphology and biochemistry of male adolescent rats

G Kerimoğlu, T Mercantepe, HS Erol, A Turgut, H Kaya, S Çolakoğlu & ...show all

Pages 445-454 | Accepted 19 Jul 2016, Published online: 11 Aug 2016

Cite this article https://doi.org/10.1080/10520295.2016.1216165 Check for updates

Sample our Bioscience Journals
>> Sign in here to start your access to the latest two volumes for 14 days

Full Article Figures & data References Citations Metrics

Reprints & Permissions Read this article

Abstract

The pathological effects of exposure to an electromagnetic field (EMF) during adolescence may be greater than those in adulthood. We investigated the effects of exposure to 900 MHz EMF during adolescence on male adult rats. Twenty-four 21-day-old male rats were divided into three equal groups: control (Cont-Gr), sham (Shm-Gr) and EMF-exposed (EMF-Gr). EMF-Gr rats were placed in an EMF exposure cage (Plexiglas

Gr rats were placed inside the Plexiglas cage under the same conditions and for the same duration, but were not exposed to EMF. All animals were sacrificed on postnatal day 60 and the hearts were extracted for microscopic and biochemical analyses. Biochemical analysis showed increased levels of malondialdehyde and superoxide dismutase, and reduced glutathione and catalase levels in EMF-Gr compared to Cont-Gr animals. Hematoxylin and eosin stained sections from EMF-Gr animals exhibited structural changes and capillary congestion in the myocardium. The percentage of apoptotic myocardial cells in EMF-Gr was higher than in either Shm-Gr or Cont-Gr animals. Transmission electron microscopy of myocardial cells of EMF-Gr animals showed altered structure of Z bands, decreased myofilaments and pronounced vacuolization. We found that exposure of male rats to 900 MHz EMF for 1 h/day during adolescence caused oxidative stress, which caused structural alteration of male adolescent rat heart tissue.

Key words:: adolescent apoptosis cellular phone electromagnetic field electron microcopy heart rat

Previous article **View** issue table of contents **Next** article

Acknowledgment

Our work was supported by the Karadeniz Technical University Research Fund, Project number: 12222.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Share

Related research